MULTIMEDIA UNIVERSITY

STUDENT ID NO

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 2, 2016/2017

BMS1824 – MANAGERIAL STATISTICS

(All sections / Groups)

28 FEBRUARY 2017
2.30 p.m – 4.30 p.m
(2 Hours)

INSTRUCTIONS TO STUDENTS

1. This question paper consists of **THIRTEEN (13)** printed pages with:
   Section A: Ten (10) multiple choice questions (20%)
   Section B: Three (3) structured questions (80%)

2. Answer **ALL** questions.

3. Answer **Section A** in the multiple-choice answer sheet provided and **Section B** in the answer booklet provided.

4. Statistical tables are attached at the end of the question paper.

5. Students are allowed to use non-programmable scientific calculators with no restrictions.

SECTION A: MULTIPLE CHOICE QUESTIONS (20 MARKS)

There are TEN (10) questions in this section. Answer ALL questions on the multiple choice answer sheet.

1. Data that is ordered or ranked according to some relationship to one another are called _____ data.
   A. ordinal
   B. ratio
   C. nominal
   D. interval

2. The probability of Derick hitting a homerun is one chance in fifty. What is the probability that he will **not hit** a homerun?
   A. 2%
   B. 98%
   C. 50%
   D. unknown since the events are mutually exclusive

3. The following data represents the crude oil prices in Malaysia per barrel for year 2007:

   RM52.11 RM54.63 RM57.83 RM63.40 RM64.93

   Determine the median price of crude oil in 2007.
   A. RM52.11
   B. RM54.63
   C. RM57.83
   D. RM63.40

4. Assume that the number of customers who arrive at a water ice stand follows the Poisson distribution with an average rate of 6.4 per hour. State the variance of this Poisson distribution.
   A. 0.5 per hour
   B. 1 per hour
   C. 3.2 per hour
   D. 6.4 per hour

5. Given that a sample of size 32 and a population standard deviation of 0.3, what would be the margin of error at a 90% level of confidence?
   A. ± 0.087
   B. ± 0.3
   C. ± 32
   D. ± 0.445

Continued...

6. If the null hypothesis is false, it is more likely that the test statistic will _____.
   A. not follow the normal distribution
   B. fall into the rejection region
   C. not follow the Poisson distribution
   D. not be symmetrical

7. According to the Health Ministry, 16% of nursing homes received five stars in 2011. A random sample of six nursing homes were selected. What is the probability that two of them received five stars?
   A. 0.2011
   B. 0.2397
   C. 0.1912
   D. 0.1684

8. The federal government would like to test the hypothesis that the average age of men filing for Social Security is higher than the average age of women set using $\alpha = 5\%$ with the following data:

	Men	Women
Sample mean	64.5 years	63.6 years
Sample size	35	39
Population standard deviation	3.0 years	3.5 years

   If Population 1 is defined as men and Population 2 is defined as women, the critical value for this hypothesis test would be _____.
   A. -1.96
   B. 2.5758
   C. 1.96
   D. 1.6449

9. What is the appropriate alternative hypothesis for the null hypothesis, 'H_0: mean production costs = RM 2,000'?
   A. H_1: mean production costs $\neq$ RM 2,000
   B. H_1: mean production costs $<$ RM 2,000
   C. H_1: mean production costs $>$ RM 2,000
   D. H_1: mean production costs $\geq$ RM 2,000

Continued...

10. The following data set represents the age (in years) of a random sample of cars on the road today.

1	3	3	7	10	12

Calculate the variance of this sample.

A. 12.9
B. 19.2
C. 4.38
D. 3.0

Continued...

SECTION B: STRUCTURED QUESTIONS (80 MARKS)

There are THREE questions in this section. Candidates MUST answer ALL THREE questions.

Question 1 (25 Marks)

(a) In a certain microwave oven on the high power setting, the time it takes for a randomly chosen kernel of popcorn to pop is normally distributed with a mean of 140 seconds and a standard deviation of 25 seconds.

(i) What is the probability of the popcorn kernel to pop less than 120 seconds?

(4 marks)

(ii) What percentage of the popcorn kernel to pop less than three minutes?

(4 marks)

(iii) What time would you allow if you wanted 95 percent of the kernels to pop?

(5 marks)

(b) The following table shows the number of car accidents reported each day in a particular city over the last 50 days:

Number of accidents	1	2	3	4	5
Frequency (days)	23	15	7	3	2

(i) What is the probability from one until two accidents occur?

(3 marks)

(ii) Determine the mean number of car accidents reported each day.

(4 marks)

(iii) Find the standard deviation for reported car accidents daily.

(5 marks)

Continued...

Question 2 (25 Marks)

(a) There were 18 depositors received some cash back from 43 bank customers who depositing a check.

(i) Construct a 90% confidence interval for the depositors who receive the cash back.

(7 marks)

(ii) How large a sample of depositors would be needed to estimate using 95 percent confidence and an error of ± 0.05?

(5 marks)

(b) Researchers analyzed 12 samples of two kinds of Stella's decaffeinated coffee. The caffeine in a cup of decaffeinated espresso had a mean of 9.4 mg with a standard deviation of 3.2 mg, while in a cup of brewed decaffeinated coffee had a mean of 12.7 mg with a standard deviation of 0.35 mg. Is there any significant difference of the caffeine content between these two beverage at $\alpha = 5\%$?

(13 marks)

Question 3 (30 Marks)

(a) Movie theaters continue to draw more people than all theme parks and major sports combined. A movie still provides the most affordable entertainment option. A statistician was observed a relationship of the moviegoers' snack spending according to the age distribution. These data, scatterplot and summary output of the relationship between the two variables are also shown below:

Age	Snack Expenses (RM)
30	2.85
50	6.5
34	1.5
12	6.35
37	6.2
33	6.75
36	3.6
26	6.1
18	8.35
46	4.35

Continued...



Summary Output

Regression Statistics	
Multiple R	0.2918
R Square	0.0851
Adjusted R Square	-0.0292
Standard Error	2.1284
Observations	10

ANOVA

	df	SS	MS	F	Significance F
Regression	1	3.373	3.373	0.745	0.413
Residual	8	36.240	4.530		
Total	9	39.612			

	Coefficients	Standard Error	t Stat	P-value
Intercept	6.9609	2.088	3.333	0.010
Age	-0.0530	0.061	-0.863	0.413

(i) Based on the scatter plot above, describe the relationship between the age distribution and the snack expenses by the moviegoers.

(3 marks)

Continued...

(ii) Determine the least square regression equation to depict the relationship between the two variables. State the unit of measurement for each variable.

(4 marks)

(iii) Interpret the slope coefficient for the age distribution which affect the snack expenses by the moviegoers.

(3 marks)

(iv) Interpret the coefficient of determination between the age distribution which affect the snack expenses by a moviegoer.

(4 marks)

(v) Predict the snack expenses if the ages are set at 10 and 45 years old. Are these estimates reliable? Explain.

(6 marks)

(b) The prices and the numbers of various items produced by a small machine and stamping plant are reported below:

Item	2000		2009	
	Price (RM)	Quantity	Price (RM)	Quantity
Washer	0.07	17 000	0.10	20 000
Cotter pin	0.04	125 000	0.03	130 000
Stove bolt	0.15	40 000	0.15	42 000
Hex nut	0.08	62 000	0.10	65 000

Compute and interpret the Laspeyres Price Index (LPI) and Paashe Price Index (PPI) for 2009 using 2000 as the base period.

(10 marks)

End of Page

A. STATISTICAL FORMULAE

A. DESCRIPTIVE STATISTICS

Mean $(\bar{x}) = \frac{\sum_{i=1}^{n} X_i}{n}$

Standard Deviation $(s) = \sqrt{\frac{\sum_{i=1}^{n} X_i^2}{n-1} - \frac{(\sum_{i=1}^{n} X_i)^2}{n(n-1)}}$

Coefficient of Variation $(CV) = \frac{\sigma}{\bar{X}} \times 100$

Pearson's Coefficient of Skewness $(S_k) = \frac{3(\bar{X} - Median)}{s}$

B. PROBABILITY

P (A or B) = P (A) + P (B) – P (A and B)

P (A and B) = P (A)× P (B) if A and B are independent

P (A | B) = P (A and B) ÷ P (B)

Poisson Probability Distribution

If X follows a Poisson Distribution , $P(\lambda)$ where $P(X = x) = \frac{e^{-\lambda}\lambda^x}{x!}$

then the mean = $E(X) = \lambda$ and variance = $VAR(X) = \lambda$

Binomial Probability Distribution

If X follows a Binomial Distribution B(n, p) where $P(X = x) = {}^nC_x p^x q^{n-x}$

then the mean = $E(X) = np$ and variance = $VAR(X) = npq$ where $q = 1 - p$

Normal Distribution

If X follows a Normal distribution, $N(\mu, \sigma)$ where $E(X) = \mu$ and $VAR(X) = \sigma^2$

then $Z = \frac{X - \mu}{\sigma}$

C. EXPECTATION AND VARIANCE OPERATORS

$E(X) = \sum[X \bullet P(X)]$

$VAR(X) = E(X^2) - [E(X)]^2$ where $E(X^2) = \sum[X^2 \bullet P(X)]$

If $E(X) = \mu$ then $E(cX) = c\,\mu$, $E(X_1 + X_2) = E(X_1) + E(X_2)$

If $VAR(X) = \sigma^2$ then $VAR(cX) = c^2\,\sigma^2$,

$VAR(X_1 + X_2) = VAR(X_1) + VAR(X_2) + 2\,COV(X_1, X_2)$

where $COV(X_1, X_2) = E(X_1 X_2) - [E(X_1)\,E(X_2)]$

D. CONFIDENCE INTERVAL ESTIMATION AND SAMPLE SIZE DETERMINATION

(100 – α) % Confidence Interval for Population Mean (σ Known) = $\mu = \overline{X} \pm Z_{\alpha/2}\left(\sigma/\sqrt{n}\right)$

(100 – α)% Confidence Interval for Population Mean (σ Unknown) = $\mu = \overline{X} \pm t_{\alpha/2,n-1}\left(s/\sqrt{n}\right)$

(100 – α)% Confidence Interval for Population Proportion $= \hat{p} \pm Z_{\alpha/2}\sigma_{p^\wedge}$

Where $\sigma_{\hat{p}} = \sqrt{\dfrac{\hat{p}(1-\hat{p})}{n}}$

Sample Size Determination for Population Mean $= n \geq \left[\dfrac{(Z_{\alpha/2})\sigma}{E}\right]^2$

Sample Size Determination for Population Proportion $= n \geq \dfrac{(Z_{\alpha/2})^2\,\hat{p}(1-\hat{p})}{E^2}$

Where E = Limit of Error in Estimation

E. HYPOTHESIS TESTING

One Sample Mean Test

Standard Deviation (σ) Known	**Standard Deviation (σ) Not Known**
$Z = \dfrac{\bar{x} - \mu}{\sigma/\sqrt{n}}$	$t = \dfrac{\bar{x} - \mu}{s/\sqrt{n}}$

One Sample Proportion Test

$z = \dfrac{\hat{p} - p}{\sigma_p}$ where $\sigma_p = \sqrt{\dfrac{p(1-p)}{n}}$

Two Sample Mean Test

Standard Deviation (σ) Known	**Standard Deviation (σ) Not Known**
$z = \dfrac{(\overline{x}_1 - \overline{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\sigma_1^2/n_1 + \sigma_2^2/n_2}}$	$t = \dfrac{(\overline{x}_1 - \overline{x}_2) - (\mu_1 - \mu_2)}{\sqrt{S_p^2\left(1/n_1 + 1/n_2\right)}}$ where $S_p^2 = \dfrac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{(n_1 + n_2 - 2)}$

Two Sample Proportion Test

$Z = \dfrac{\hat{p}_1 - \hat{p}_2}{\sqrt{p(1-p)\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$ where $p = \dfrac{X_1 + X_2}{n_1 + n_2}$

where X_1 and X_2 are the number of successes from each population

F. REGRESSION ANALYSIS

Simple Linear Regression

Population Model: $Y = \beta_0 + \beta_1 X_1 + \varepsilon$

Sample Model: $\hat{y} = b_0 + b_1 x_1 + e$

Correlation Coefficient

$$r = \frac{\sum XY - \left[\frac{\sum X \sum Y}{n}\right]}{\sqrt{\left[\sum X^2 - \left((\sum X)^2 / n\right)\right]\left[\sum Y^2 - \left((\sum Y)^2 / n\right)\right]}} = \frac{COV(X,Y)}{\sigma_X \sigma_Y}$$

ANOVA Table for Regression

Source	Degrees of Freedom	Sum of Squares	Mean Squares
Regression	*1*	SSR	MSR = SSR/1
Error/Residual	$n-2$	SSE	MSE = SSE/$(n-2)$
Total	$n-1$	SST	

Test Statistic for Significance of the Predictor Variable

$t_i = \frac{b_i}{S_{b_i}}$ and the critical value $= \pm t_{\alpha/2,(n-p-1)}$

Where p= number of predictor

G. INDEX NUMBERS

Simple Price Index

$$P = \frac{p_t}{p_0} \times 100$$

Laspeyres Quantity Index

$$P = \frac{\sum p_0 q_t}{\sum p_0 q_0} \times 100$$

Aggregate Price Index

$$P = \frac{\sum p_t}{\sum p_0}(100)$$

Paasche Quantity Index

$$P = \frac{\sum p_t q_t}{\sum p_t q_0} \times 100$$

Laspeyres Price Index

$$P = \frac{\sum p_t q_0}{\sum p_0 q_0} \times 100$$

Fisher's Ideal Price Index

$$\sqrt{(Laspeyres\ \ Price\ Index)(Paa\ sche\ Price\ Index)}$$

Paasche Price Index

$$P = \frac{\sum p_t q_t}{\sum p_0 q_t} \times 100$$

Value Index

$$V = \frac{\sum p_t q_t}{\sum p_0 q_0} \times 100$$

B. STATISTICAL TABLE

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

The function tabulated is $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{1}{2}t^2}\, dt$. $\Phi(x)$ is the probability that a random variable, normally distributed with zero mean and unit variance, will be less than or equal to x. When $x < 0$ use $\Phi(x) = 1 - \Phi(-x)$, as the normal distribution with zero mean and unit variance is symmetric about zero.



x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0·00	0·5000	0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·97725
·01	·5040	·41	·6591	·81	·7910	·21	·8869	·61	·9463	·01	·97778
·02	·5080	·42	·6628	·82	·7939	·22	·8888	·62	·9474	·02	·97831
·03	·5120	·43	·6664	·83	·7967	·23	·8907	·63	·9484	·03	·97882
·04	·5160	·44	·6700	·84	·7995	·24	·8925	·64	·9495	·04	·97932
0·05	0·5199	0·45	0·6736	0·85	0·8023	1·25	0·8944	1·65	0·9505	2·05	0·97982
·06	·5239	·46	·6772	·86	·8051	·26	·8962	·66	·9515	·06	·98030
·07	·5279	·47	·6808	·87	·8078	·27	·8980	·67	·9525	·07	·98077
·08	·5319	·48	·6844	·88	·8106	·28	·8997	·68	·9535	·08	·98124
·09	·5359	·49	·6879	·89	·8133	·29	·9015	·69	·9545	·09	·98169
0·10	0·5398	0·50	0·6915	0·90	0·8159	1·30	0·9032	1·70	0·9554	2·10	0·98214
·11	·5438	·51	·6950	·91	·8186	·31	·9049	·71	·9564	·11	·98257
·12	·5478	·52	·6985	·92	·8212	·32	·9066	·72	·9573	·12	·98300
·13	·5517	·53	·7019	·93	·8238	·33	·9082	·73	·9582	·13	·98341
·14	·5557	·54	·7054	·94	·8264	·34	·9099	·74	·9591	·14	·98382
0·15	0·5596	0·55	0·7088	0·95	0·8289	1·35	0·9115	1·75	0·9599	2·15	0·98422
·16	·5636	·56	·7123	·96	·8315	·36	·9131	·76	·9608	·16	·98461
·17	·5675	·57	·7157	·97	·8340	·37	·9147	·77	·9616	·17	·98500
·18	·5714	·58	·7190	·98	·8365	·38	·9162	·78	·9625	·18	·98537
·19	·5753	·59	·7224	·99	·8389	·39	·9177	·79	·9633	·19	·98574
0·20	0·5793	0·60	0·7257	1·00	0·8413	1·40	0·9192	1·80	0·9641	2·20	0·98610
·21	·5832	·61	·7291	·01	·8438	·41	·9207	·81	·9649	·21	·98645
·22	·5871	·62	·7324	·02	·8461	·42	·9222	·82	·9656	·22	·98679
·23	·5910	·63	·7357	·03	·8485	·43	·9236	·83	·9664	·23	·98713
·24	·5948	·64	·7389	·04	·8508	·44	·9251	·84	·9671	·24	·98745
0·25	0·5987	0·65	0·7422	1·05	0·8531	1·45	0·9265	1·85	0·9678	2·25	0·98778
·26	·6026	·66	·7454	·06	·8554	·46	·9279	·86	·9686	·26	·98809
·27	·6064	·67	·7486	·07	·8577	·47	·9292	·87	·9693	·27	·98840
·28	·6103	·68	·7517	·08	·8599	·48	·9306	·88	·9699	·28	·98870
·29	·6141	·69	·7549	·09	·8621	·49	·9319	·89	·9706	·29	·98899
0·30	0·6179	0·70	0·7580	1·10	0·8643	1·50	0·9332	1·90	0·9713	2·30	0·98928
·31	·6217	·71	·7611	·11	·8665	·51	·9345	·91	·9719	·31	·98956
·32	·6255	·72	·7642	·12	·8686	·52	·9357	·92	·9726	·32	·98983
·33	·6293	·73	·7673	·13	·8708	·53	·9370	·93	·9732	·33	·99010
·34	·6331	·74	·7704	·14	·8729	·54	·9382	·94	·9738	·34	·99036
0·35	0·6368	0·75	0·7734	1·15	0·8749	1·55	0·9394	1·95	0·9744	2·35	0·99061
·36	·6406	·76	·7764	·16	·8770	·56	·9406	·96	·9750	·36	·99086
·37	·6443	·77	·7794	·17	·8790	·57	·9418	·97	·9756	·37	·99111
·38	·6480	·78	·7823	·18	·8810	·58	·9429	·98	·9761	·38	·99134
·39	·6517	·79	·7852	·19	·8830	·59	·9441	·99	·9767	·39	·99158
0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·9772	2·40	0·99180

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
2·40	0·99180	2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918
·41	·99202	·56	·99477	·71	·99664	·86	·99788	·01	·99869	·16	·99921
·42	·99224	·57	·99492	·72	·99674	·87	·99795	·02	·99874	·17	·99924
·43	·99245	·58	·99506	·73	·99683	·88	·99801	·03	·99878	·18	·99926
·44	·99266	·59	·99520	·74	·99693	·89	·99807	·04	·99882	·19	·99929
2·45	0·99286	2·60	0·99534	2·75	0·99702	2·90	0·99813	3·05	0·99886	3·20	0·99931
·46	·99305	·61	·99547	·76	·99711	·91	·99819	·06	·99889	·21	·99934
·47	·99324	·62	·99560	·77	·99720	·92	·99825	·07	·99893	·22	·99936
·48	·99343	·63	·99573	·78	·99728	·93	·99831	·08	·99896	·23	·99938
·49	·99361	·64	·99585	·79	·99736	·94	·99836	·09	·99900	·24	·99940
2·50	0·99379	2·65	0·99598	2·80	0·99744	2·95	0·99841	3·10	0·99903	3·25	0·99942
·51	·99396	·66	·99609	·81	·99752	·96	·99846	·11	·99906	·26	·99944
·52	·99413	·67	·99621	·82	·99760	·97	·99851	·12	·99910	·27	·99946
·53	·99430	·68	·99632	·83	·99767	·98	·99856	·13	·99913	·28	·99948
·54	·99446	·69	·99643	·84	·99774	·99	·99861	·14	·99916	·29	·99950
2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918	3·30	0·99952

The critical table below gives on the left the range of values of x for which $\Phi(x)$ takes the value on the right, correct to the last figure given; in critical cases, take the upper of the two values of $\Phi(x)$ indicated.

3·075	0·9990	3·263	0·9994	3·731	0·99990	3·916	0·99995
3·105	0·9991	3·320	0·9995	3·759	0·99991	3·976	0·99996
3·138	0·9992	3·389	0·9996	3·791	0·99992	4·055	0·99997
3·174	0·9993	3·480	0·9997	3·826	0·99993	4·173	0·99998
3·215	0·9994	3·615	0·9998	3·867	0·99994	4·417	0·99999
			0·9999		0·99995		1·00000

When $x > 3{\cdot}3$ the formula $1-\Phi(x) \doteq \frac{e^{-\frac{1}{2}x^2}}{x\sqrt{2\pi}}\left[1-\frac{1}{x^2}+\frac{3}{x^4}-\frac{15}{x^6}+\frac{105}{x^8}\right]$ is very accurate, with relative error less than $945/x^{10}$.

TABLE 5. PERCENTAGE POINTS OF THE NORMAL DISTRIBUTION

This table gives percentage points $x(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{2\pi}}\int_{x(P)}^{\infty} e^{-\frac{1}{2}t^2}\,dt.$$

If X is a variable, normally distributed with zero mean and unit variance, $P/100$ is the probability that $X \geqslant x(P)$. The lower P per cent points are given by symmetry as $-x(P)$, and the probability that $|X| \geqslant x(P)$ is $2P/100$.



P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$
50	0·0000	5·0	1·6449	3·0	1·8808	2·0	2·0537	1·0	2·3263	0·10	3·0902
45	0·1257	4·8	1·6646	2·9	1·8957	1·9	2·0749	0·9	2·3656	0·09	3·1214
40	0·2533	4·6	1·6849	2·8	1·9110	1·8	2·0969	0·8	2·4089	0·08	3·1559
35	0·3853	4·4	1·7060	2·7	1·9268	1·7	2·1201	0·7	2·4573	0·07	3·1947
30	0·5244	4·2	1·7279	2·6	1·9431	1·6	2·1444	0·6	2·5121	0·06	3·2389
25	0·6745	4·0	1·7507	2·5	1·9600	1·5	2·1701	0·5	2·5758	0·05	3·2905
20	0·8416	3·8	1·7744	2·4	1·9774	1·4	2·1973	0·4	2·6521	0·01	3·7190
15	1·0364	3·6	1·7991	2·3	1·9954	1·3	2·2262	0·3	2·7478	0·005	3·8906
10	1·2816	3·4	1·8250	2·2	2·0141	1·2	2·2571	0·2	2·8782	0·001	4·2649
5	1·6449	3·2	1·8522	2·1	2·0335	1·1	2·2904	0·1	3·0902	0·0005	4·4172

TABLE 10. PERCENTAGE POINTS OF THE t-DISTRIBUTION

This table gives percentage points $t_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{\nu\pi}} \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2})}{\Gamma(\frac{1}{2}\nu)} \int_{t_\nu(P)}^{\infty} \frac{dt}{(1+t^2/\nu)^{\frac{1}{2}(\nu+1)}}.$$

Let X_1 and X_2 be independent random variables having a normal distribution with zero mean and unit variance and a χ^2-distribution with ν degrees of freedom respectively; then $t = X_1/\sqrt{X_2/\nu}$ has Student's t-distribution with ν degrees of freedom, and the probability that $t \geqslant t_\nu(P)$ is $P/100$. The lower percentage points are given by symmetry as $-t_\nu(P)$, and the probability that $|t| \geqslant t_\nu(P)$ is $2P/100$.



The limiting distribution of t as ν tends to infinity is the normal distribution with zero mean and unit variance. When ν is large interpolation in ν should be harmonic.

P	40	30	25	20	15	10	5	2·5	1	0·5	0·1	0·05
$\nu = 1$	0·3249	0·7265	1·0000	1·3764	1·963	3·078	6·314	12·71	31·82	63·66	318·3	636·6
2	0·2887	0·6172	0·8165	1·0607	1·386	1·886	2·920	4·303	6·965	9·925	22·33	31·60
3	0·2767	0·5844	0·7649	0·9785	1·250	1·638	2·353	3·182	4·541	5·841	10·21	12·92
4	0·2707	0·5686	0·7407	0·9410	1·190	1·533	2·132	2·776	3·747	4·604	7·173	8·610
5	0·2672	0·5594	0·7267	0·9195	1·156	1·476	2·015	2·571	3·365	4·032	5·893	6·869
6	0·2648	0·5534	0·7176	0·9057	1·134	1·440	1·943	2·447	3·143	3·707	5·208	5·959
7	0·2632	0·5491	0·7111	0·8960	1·119	1·415	1·895	2·365	2·998	3·499	4·785	5·408
8	0·2619	0·5459	0·7064	0·8889	1·108	1·397	1·860	2·306	2·896	3·355	4·501	5·041
9	0·2610	0·5435	0·7027	0·8834	1·100	1·383	1·833	2·262	2·821	3·250	4·297	4·781
10	0·2602	0·5415	0·6998	0·8791	1·093	1·372	1·812	2·228	2·764	3·169	4·144	4·587
11	0·2596	0·5399	0·6974	0·8755	1·088	1·363	1·796	2·201	2·718	3·106	4·025	4·437
12	0·2590	0·5386	0·6955	0·8726	1·083	1·356	1·782	2·179	2·681	3·055	3·930	4·318
13	0·2586	0·5375	0·6938	0·8702	1·079	1·350	1·771	2·160	2·650	3·012	3·852	4·221
14	0·2582	0·5366	0·6924	0·8681	1·076	1·345	1·761	2·145	2·624	2·977	3·787	4·140
15	0·2579	0·5357	0·6912	0·8662	1·074	1·341	1·753	2·131	2·602	2·947	3·733	4·073
16	0·2576	0·5350	0·6901	0·8647	1·071	1·337	1·746	2·120	2·583	2·921	3·686	4·015
17	0·2573	0·5344	0·6892	0·8633	1·069	1·333	1·740	2·110	2·567	2·898	3·646	3·965
18	0·2571	0·5338	0·6884	0·8620	1·067	1·330	1·734	2·101	2·552	2·878	3·610	3·922
19	0·2569	0·5333	0·6876	0·8610	1·066	1·328	1·729	2·093	2·539	2·861	3·579	3·883
20	0·2567	0·5329	0·6870	0·8600	1·064	1·325	1·725	2·086	2·528	2·845	3·552	3·850
21	0·2566	0·5325	0·6864	0·8591	1·063	1·323	1·721	2·080	2·518	2·831	3·527	3·819
22	0·2564	0·5321	0·6858	0·8583	1·061	1·321	1·717	2·074	2·508	2·819	3·505	3·792
23	0·2563	0·5317	0·6853	0·8575	1·060	1·319	1·714	2·069	2·500	2·807	3·485	3·768
24	0·2562	0·5314	0·6848	0·8569	1·059	1·318	1·711	2·064	2·492	2·797	3·467	3·745
25	0·2561	0·5312	0·6844	0·8562	1·058	1·316	1·708	2·060	2·485	2·787	3·450	3·725
26	0·2560	0·5309	0·6840	0·8557	1·058	1·315	1·706	2·056	2·479	2·779	3·435	3·707
27	0·2559	0·5306	0·6837	0·8551	1·057	1·314	1·703	2·052	2·473	2·771	3·421	3·690
28	0·2558	0·5304	0·6834	0·8546	1·056	1·313	1·701	2·048	2·467	2·763	3·408	3·674
29	0·2557	0·5302	0·6830	0·8542	1·055	1·311	1·699	2·045	2·462	2·756	3·396	3·659
30	0·2556	0·5300	0·6828	0·8538	1·055	1·310	1·697	2·042	2·457	2·750	3·385	3·646
32	0·2555	0·5297	0·6822	0·8530	1·054	1·309	1·694	2·037	2·449	2·738	3·365	3·622
34	0·2553	0·5294	0·6818	0·8523	1·052	1·307	1·691	2·032	2·441	2·728	3·348	3·601
36	0·2552	0·5291	0·6814	0·8517	1·052	1·306	1·688	2·028	2·434	2·719	3·333	3·582
38	0·2551	0·5288	0·6810	0·8512	1·051	1·304	1·686	2·024	2·429	2·712	3·319	3·566
40	0·2550	0·5286	0·6807	0·8507	1·050	1·303	1·684	2·021	2·423	2·704	3·307	3·551
50	0·2547	0·5278	0·6794	0·8489	1·047	1·299	1·676	2·009	2·403	2·678	3·261	3·496
60	0·2545	0·5272	0·6786	0·8477	1·045	1·296	1·671	2·000	2·390	2·660	3·232	3·460
120	0·2539	0·5258	0·6765	0·8446	1·041	1·289	1·658	1·980	2·358	2·617	3·160	3·373
∞	0·2533	0·5244	0·6745	0·8416	1·036	1·282	1·645	1·960	2·326	2·576	3·090	3·291